

# 5-CHROM-44 UROKINASE CHROMOGENIC SUBSTRATE REF 5D-30806

For Research Use Only.

Not for Use in Diagnostic Procedures.

Store at 2-8°C 25 mg

### ORIGIN

Synthetic generic Chromogenic substrate, highly purified and stabilized.

#### **SPECIFICITY**

Recommended substrate for Urokinase.

#### **PRESENTATION**

Vial containing 25 mg of 5-CHROM-44, lyophilized in presence of Mannitol as a bulking agent.

#### RECONSTITUTION

According to the research protocol used, the 5-CHROM-44 chromogenic substrate can be restored with variable volumes of distilled water; for example, 5 mL can be used for a substrate concentration of 5 mg/mL, or 20 mL for a substrate concentration of 1.25 mg/mL.

## PEPTIDE SEQUENCE

pyro-Glu-Gly-Arg-pNa.HCl

# **CHEMICAL STRUCTURE**

C<sub>19</sub>H<sub>26</sub>N<sub>8</sub>O<sub>6</sub>, HCl

## **MOLARITY**

≈54 µmol/vial

## **PURITY GRADE**

> 95 %

# **MOLECULAR WEIGHT**

462.5

# FREE pNA CONTENT (MEASURED ON SUBSTRATE AT 2.5 g/L)

 $OD_{405nm} \le 0.300$ 

## USE

All research studies and protocols where a source of chromogenic substrate for Urokinase is required. This kit is for research use only and should not be used for patient diagnosis or treatment.

## RESPECTIVE REACTIVITIES

APC	FXa	Plasmin	Kallikrein	Thrombin	Urokinase
1	2	1	2	<1	100

Assay conditions must be established for making the substrate totally specific for Urokinase.

## **STABILITY**

## Restored:

- 7 days at room temperature
- 3 months at 2-8°C
- Do not freeze

# Lyophilized:

• Until the expiration date printed on the vial.



5-Diagnostics AG

Heuberg 7, 4051 Basel, Switzerland www.5-diagnostics.com info@5-diagnostics.com